

Listeria Fraser Broth

Liquid medium for the detection of *Listeria monocytogenes* and *Listeria* spp, according to ISO 11290-1.

TYPICAL FORMULA	(g/l)
Enzymatic Digest of Animal Tissues	5.0
Enzymatic Digest of Casein	5.0
Meat Extract	5.0
Yeast Extract	5.0
Sodium Chloride	20.0
Disodium Phosphate, Anhydrous	9.6*
Potassium Dihydrogen Phosphate	1.35
Aesculin	1.0
Lithium Chloride	3.0
Nalidixic Acid	0.02
Acriflavine	0.025
Final pH 7.2 ± 0.2 at 25°C	

*Equivalent to 12.0 g of Disodium Hydrogen Phosphate, Dihydrate.

DESCRIPTION

Demi Fraser Broth is a liquid medium used with supplements for the secondary enrichment of *L. monocytogenes* and *Listeria* spp from food, animal feeding and environmental samples in the area of food production and food handling.

The medium is completed after the addition of Listeria Fraser Supplement which is available in the following forms:

- Freeze-dried (ref. 81028)
- Liquid (ref. 80304)

Both items contain Ammonium Iron(III) Citrate.

PRINCIPLE

Enzymatic digest of animal tissues, enzymatic digest of casein and meat extract provide nitrogen, vitamins, minerals and amino acids for organisms growth. Yeast extract is a source of vitamins, particularly of B-group. Sodium chloride maintains the osmotic balance of the medium and in a so high concentration inhibits enterococci. Potassium and sodium phosphates act as buffer system. Aesculin is hydrolyzed by all *Listeria* species to aesculetin. Lithium chloride is inhibitory for the accompanying flora. Acriflavine and nalidixic acid are selective agents.

Ferric ions provided by ammonium iron(III) citrate will react with aesculetin producing a blackening of the medium.

PREPARATION

Suspend 55 g of powder in 1 liter of deionized or distilled water. Bring to boil and shake until completely dissolved. Sterilize at 121°C for 15 minutes. Cool up to 45-50°C. Aseptically, add the content of 2 vials of Listeria Fraser Supplement. NOTE: Either ref. 81028 or ref. 80304 can be chosen. The latter is a ready to use liquid solution whereas ref. 81028 needs to be reconstituted with water before use.

TECHNIQUE

1. Add sample to Demi Fraser Broth (ref. 452401) to prepare a 10-fold dilution (w/v or v/v).
Incubate at 30 ± 1°C for 25 ± 1 h.
2. Transfer 0.1 ml of the primary enrichment culture into 10 ml of Listeria Fraser Broth.
Incubate at 37 ± 1°C for 24 ± 2 h.
3. Streak from both primary and secondary enrichments onto O.A. Listeria Agar (ref. 10620) to obtain well-separated colonies.
Incubate at 37 ± 1°C for 24 ± 2 h and for an additional 24 ± 2 h.
4. Use the selective enrichments to inoculate a second selective medium, e.g. Listeria Palcam Agar (ref. 10041), Listeria Oxford Agar (ref. 61016). Refer to the relevant technical sheet for further details.

INTERPRETATION OF RESULTS

A blackening of Demi Fraser Broth can be observed after incubation.

Blue-green colonies with or without halo on O.A. Listeria Agar are considered presumptive *Listeria* spp. Typical colonies of *L. monocytogenes* are surrounded by an opaque halo.

For confirmation, subculture onto appropriate non-selective agar, e.g. Blood Agar, Nutrient Agar, TSYEA (ref. 10432). Then, carry out confirmation tests including a positive and negative control.

STORAGE CONDITIONS

The powder is very hygroscopic, store the powder at 10-30°C, in a dry environment, in its original container tightly closed and use it before the expiry date on the label or until signs of deterioration or contamination are evident. Store prepared medium at 2-8°C away from light.

WARNING AND PRECAUTIONS

The product does not contain hazardous substances in concentrations exceeding the limits set by current legislation and therefore is not classified as dangerous. It is nevertheless recommended to consult the safety data sheet for its correct use. The product is designed for professional use only and must be used by properly trained operators.

DISPOSAL OF WASTE

Disposal of waste must be carried out according to the national and local regulations in force.



LIOFILCHEM® S.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto degli Abruzzi (TE) - ITALY
Tel +39 0858930745 Fax +39 0858930330 www.liofilchem.com

REFERENCES

1. ISO 11290-1:2017. Microbiology of the food chain – Horizontal method for the detection and enumeration of *Listeria monocytogenes* and *Listeria* spp. – Part 1: Detection Method.
2. ISO 11290-2:2017. Microbiology of the food chain – Horizontal method for the detection and enumeration of *Listeria monocytogenes* and *Listeria* spp. – Part 2: Enumeration Method.
3. EN ISO 11133:2014. Microbiology of food, animal feed and water – Preparation, production, storage and performance testing of culture media.
4. Rapporto ISTISAN 96/35. ISSN 1123-3117. Metodi di analisi per il controllo microbiologico degli alimenti.
5. Normalisation Francaise, AFNOR (1993) V08-55.
6. Fraser. J.A and Sperber W.H (1988) J. Food Prot , 51, 762-765.



PRODUCT SPECIFICATIONS

NAME

Listeria Fraser Broth

PRESENTATION

Dehydrated medium

STORAGE

10-30°C

PACKAGING

Ref.	Content	Packaging
610166	500 g	500 g of powder in plastic bottle
620166	100 g	100 g of powder in plastic bottle

pH OF THE MEDIUM

7.2 ± 0.2

USE

Listeria Fraser Broth is a liquid medium used with supplements for the detection of *L. monocytogenes* and *Listeria* spp in food and environmental samples, according to ISO 11290-1

TECHNIQUE

Refer to technical sheet of the product

APPEARANCE OF THE MEDIUM

Powder medium

Appearance: free-flowing, homogeneous

Colour: beige

Ready-to-use medium

Appearance: clear

Colour: amber

SHELF LIFE

4 years

QUALITY CONTROL

- Control of general characteristics, label and print
- Microbiological control
Incubation Conditions: 24 ± 2 h / 37 ± 1°C

Inoculum for productivity: ≤100 CFU











Microorganism	Specification
<i>Listeria monocytogenes</i> serovar 4b	WDCM 00021
+ <i>Escherichia coli</i>	WDCM 00013
+ <i>Enterococcus faecalis</i>	WDCM 00009
	Blackening of the medium, >10 colonies on O.A. Listeria Agar

Inoculum for selectivity: >10³ CFU

Microorganism	Specification
<i>Escherichia coli</i>	WDCM 00013
<i>Enterococcus faecalis</i>	WDCM 00009
	Total inhibition on TSA
	<100 colonies on TSA

TSA: Tryptic Soy Agar

TABLE OF SYMBOLS

 LOT	Batch code	 Keep away from Sunlight	 Manufacturer	 Use by	 Fragile, handle with care
 REF	Catalogue number	 Temperature limitation	 Contains sufficient for <n> tests	 Caution, consult instructions for use	 Do not reuse



LIOFILCHEM® S.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto degli Abruzzi (TE) - ITALY
Tel +39 0858930745 Fax +39 0858930330 www.liofilchem.com